

Edge solitons in the QHE*

M. Hassaïne, P. A. Horváthy and J.-C. Yéra

Laboratoire de Mathématiques et de Physique Théorique
Université de Tours
Parc de Grandmont, F-37200 TOURS (France)
e-mail: horvathy@univ-tours.fr

February 1, 2008

Abstract

The spacelike reduction of the Chern-Simons Lagrangian yields a modified Nonlinear Schrödinger Equation (jNLS) where in the non-linearity the particle density is replaced by the current. When the phase is linear in the position, this latter is an ordinary NLS with time-dependent coefficients which admits interesting solutions, whose arising is explained by the conformal properties of non-relativistic spacetime. Only the usual travelling soliton is consistent with the jNLS but adding a six-order potential converts it into an integrable equation.

1 Reduction of Chern-Simons

The Landau-Ginzburg theory of the Quantum Hall Effect [1] uses the Chern-Simons Lagrangian in $(2 + 1)$ dimensions,

$$\mathcal{L} = \frac{1}{4\kappa} \epsilon_{\mu\nu\rho} F_{\mu\nu} A_\rho + i\phi^* D_t \phi - \frac{1}{2} |\vec{D}\phi|^2 - V(\phi), \quad (1)$$

where the scalar field ϕ is the order parameter and A_μ is the statistical gauge field; $D_\mu = \partial_\mu - iA_\mu$ is the covariant derivative. The constant κ is interpreted as the Hall conductivity. The second-order field equations are not integrable [2]; they admit integrable reductions, though. The simplest of these is when time-dependence is eliminated; then, for a judicious choice of the self-interaction potential $V(\phi)$, the system admits finite-energy vortex solutions [3]. Here we focus our attention to another, space-like reduction [4]. Assuming independence from one spacelike coordinate and adding a suitable kinetic term yields in fact, after elimination of the gauge field using its equation of motion,

$$\mathcal{L} = i\phi^* \partial_t \phi - \frac{1}{2} |(\partial_x - i\kappa^2 \rho)\phi|^2 - V, \quad (2)$$

where $\rho = |\phi|^2$ is the particle density. This is the model proposed in Ref. [5] to describe the edge states in the QHE.

The field equations associated to (2) read

$$\begin{aligned} i\partial_t \phi &= -\frac{1}{2}(\partial_x - i\kappa^2 \rho)^2 \phi - \kappa^2 j \phi + \frac{\partial V}{\partial \phi^*}, \\ j &= \frac{1}{2i} [\phi^* (\partial_x - i\kappa^2 \rho) \phi - \phi (\partial_x + i\kappa^2 \rho) \phi^*]. \end{aligned} \quad (3)$$

*Talk given by P. Horváthy at the meeting NONLINEARITY, INTEGRABILITY AND ALL THAT Gallipoli (Lecce) Italy, 1999. Ed. M. Boiti, L. Martina, F. Pempinelli, B. Prinari & G. Soliani, p. 474. Singapore : World Scientific (2000).

Then the particle density and the current satisfy the continuity equation $\partial_t \rho + \partial_x j = 0$. Let us first assume that $V = 0$. Now the non-local transformation [6]

$$\psi = \left(\exp[-i\kappa^2 \int^x \rho(y) dy] \right) \phi \quad (4)$$

takes (3) into the modified non-linear Schrödinger equation in which the density in the non-linearity has been replaced by the current,

$$\begin{aligned} i\partial_t \psi &= -\frac{1}{2}\partial_x^2 \psi - 2\kappa^2 j \psi, \\ j &= \frac{1}{2i}[\psi^* \partial_x \psi - \psi(\partial_x \psi)^*]. \end{aligned} \quad (5)$$

2 The variable-coefficient NLS

Decomposing ψ into module and phase, $\psi = \sqrt{\rho} e^{i\theta}$, yields (formally) the ordinary cubic NLS with variable coefficient,

$$i\partial_t \psi = -\frac{1}{2}\partial_x^2 \psi - F(t, x)|\psi|^2 \psi, \quad (6)$$

with $F(t, x) = 2\kappa^2 \partial_x \theta$. Then Aglietti et al. [4] observe that, for $\theta = vx - \omega t$, Eq. (6) reduces to the usual non-linear Schrödinger equation with constant coefficient $F = 2\kappa^2 v$ which admits, for example, the travelling soliton solution

$$\psi_s = \pm e^{i(vx - \omega t)} \sqrt{\frac{1}{2\kappa^2 v}} \frac{\alpha}{\cosh \alpha(x - vt)}, \quad \alpha^2 = v^2 - 2\omega. \quad (7)$$

The non-linearity in (6) has to be attractive, $F > 0$; the solution (7) is therefore chiral, $v > 0$.

It is natural to ask whether the travelling soliton (7) can be generalized. Let us first study the variable-coefficient NLS (6) on its own. It has been shown [7] that this equation only passes the Painlevé test of Weiss, Tabor and Carnevale [8], when the coefficient of the non-linearity is

$$F = (a + bt)^{-1}, \quad (8)$$

where a and b constants. For $b = 0$, $F(t, x)$ is a constant and we recover the constant-coefficient NLS. For $b \neq 0$, the equation becomes explicitly time-dependent. Assuming, for simplicity, that $a = 0$ and $b = 1$, it reads

$$i\partial_t \psi + \frac{1}{2}\partial_x^2 \psi + \frac{1}{t}|\psi|^2 \psi = 0. \quad (9)$$

This equation can also be solved. Generalizing the usual travelling soliton, we find, for example, the 1-soliton

$$\psi_0(t, x) = \frac{1}{\sqrt{t}} \frac{e^{i(x^2/4t - 1/2t)}}{\cosh[-x/t - x_0]}. \quad (10)$$

It is worth pointing out that the steps followed in constructing (10) are essentially the same as those for the travelling soliton of the ordinary NLS — and this is not a pure coincidence. A short calculation shows in fact that

$$\psi(t, x) = \frac{1}{\sqrt{t}} \exp\left[\frac{ix^2}{4t}\right] \Psi(-1/t, -x/t) \quad (11)$$

satisfies the time-dependent equation (9) if and only if $\Psi(t, x)$ solves Eqn. (6) with $F = 1$. Our soliton (10) comes in particular from the “standing soliton” $\Psi_s = \exp(it/2)(\cosh[x - x_0])^{-1}$ solution of the NLS.

3 Non-relativistic conformal transformations

Where does the formula (10) come from ? The non-linear space-time transformation

$$D : \begin{pmatrix} t \\ x \end{pmatrix} \rightarrow \begin{pmatrix} -1/t \\ -x/t \end{pmatrix} \quad (12)$$

has already been met in a rather different context, namely in describing planetary motion when the gravitational “constant” changes inversely with time, as suggested by Dirac [9]. One shows in fact that $\vec{r}(t) = t \vec{r}^*(-1/t)$ describes planetary motion with Newton’s “constant” varying as $G(t) = G_0/t$, whenever $\vec{r}^*(t)$ describes ordinary planetary motion, i.e. the one with a constant gravitational constant G_0 [10], [11]. The strange-looking transformation (12) is indeed related to the conformal structure of non-relativistic space-time [12]. It has been noticed in fact almost thirty years ago, that the “conformal” space-time transformations

$$\begin{aligned} \begin{pmatrix} t \\ x \end{pmatrix} &\rightarrow \begin{pmatrix} T \\ X \end{pmatrix} = \begin{pmatrix} \delta^2 t \\ \delta x \end{pmatrix}, & 0 \neq \delta \in \mathbf{R} & \text{dilatations} \\ \begin{pmatrix} t \\ x \end{pmatrix} &\rightarrow \begin{pmatrix} T \\ X \end{pmatrix} = \begin{pmatrix} \frac{t}{1 - \kappa t} \\ \frac{x}{-\kappa t} \end{pmatrix}, & \kappa \in \mathbf{R} & \text{expansions} \\ \begin{pmatrix} t \\ x \end{pmatrix} &\rightarrow \begin{pmatrix} T \\ X \end{pmatrix} = \begin{pmatrix} t + \epsilon \\ x \end{pmatrix}, & \epsilon \in \mathbf{R} & \text{time translations} \end{aligned} \quad (13)$$

implemented on wave functions according to

$$\Psi(T, X) = \begin{cases} \delta^{1/2} u(t, x) \\ (1 - \kappa t)^{1/2} \exp \left[i \frac{\kappa x^2}{4(1 - \kappa t)} \right] \psi(t, x) \\ \psi(t, x) \end{cases} \quad (14)$$

permute the solutions of the free Schrödinger equation. In other words, they are symmetries of the free Schrödinger equation. The generators in (13) span in fact an $SL(2, \mathbf{R})$ group. (A Dirac monopole, an Aharonov-Bohm vector potential and an inverse-square potential can also be included). The transformation D in Eqn. (12) belongs to this symmetry group: it is in fact (i) a time translation with $\epsilon = 1$, (ii) followed by an expansion with $\kappa = 1$, (iii) followed by a second time-translation with $\epsilon = 1$. It is hence a symmetry for the free (linear) Schrödinger equation.

The cubic NLS with constant non-linearity is not more $SL(2, \mathbf{R})$ invariant: the transformation D in (12) implemented as in Eq. (14) carries the cubic term into the time-dependent term $(1/t)|u|^2 u$, just like Newton’s gravitational potential G_0/r with $G_0 = \text{const.}$ is carried into the time-dependent Dirac expression $t^{-1}G_0/r$. More generally, the the non-linear Schrödinger equation with time-dependent coefficients,

$$i\partial_t \psi + p(t)\partial_x^2 \psi + F(t)|u|^2 \psi = 0, \quad (15)$$

can be transformed into the constant-coefficient form, whenever

$$p(t) = F(t) \left(a + b \int^t p(s) ds \right) \quad (16)$$

This same condition was found later as the one needed for the Painlevé test [7].

Similar arguments explain the integrability of other NLS type equations. For example, electromagnetic waves in a non-uniform medium propagate according to

$$i\partial_t\psi + \partial_x^2\psi + (-2\alpha x + 2|\psi|^2)\psi = 0, \quad (17)$$

which can again be solved by inverse scattering [13]. This is explained by observing that the potential term here can be eliminated by switching to a uniformly accelerated frame:

$$\begin{aligned} \psi(t, x) &= \exp[-i(2\alpha x t + \frac{4}{3}\alpha^2 t^3)]\Psi(T, X), \\ T &= t, \quad X = x + 2\alpha t^2. \end{aligned} \quad (18)$$

Then $u(t, x)$ solves (17) whenever $\Psi(T, X)$ solves the free equation.

The transformation (18) is again related to the structure of non-relativistic space-time. It can be shown in fact [11] that the (linear) Schrödinger equation

$$i\partial_t\psi + \partial_x^2\psi - V(t, x)\psi = 0 \quad (19)$$

can be brought into the free form by a space-time transformation if and only if the potential is $V(t, x) = \alpha(t)x \pm \frac{\omega^2(t)}{4}x^2$. For the uniform force field ($\omega = 0$) the required transformation is precisely (18).

For the oscillator potential ($\alpha = 0$), one can use rather Niederer's transformation [14]

$$\begin{aligned} \psi(t, x) &= (\cos \omega t)^{-1/2} \exp[-i\frac{\omega}{4}x^2 \tan \omega t] \Psi(T, X), \\ T &= \frac{\tan \omega t}{\omega} \quad X = \frac{x}{\cos \omega t}. \end{aligned} \quad (20)$$

Then

$$i\partial_t\psi + \frac{1}{2}\partial_x^2\psi - \frac{\omega^2 x^2}{4}\psi = (\cos \omega t)^{-5/2} \exp[-i\frac{\omega}{4} \tan \omega t] \left(i\partial_T\Psi + \frac{1}{2}\partial_X^2\Psi \right)$$

so that ψ satisfies the oscillator-equations iff Ψ solves the free equation.

The Niederer transformation (20) leaves the inverse square potential invariant; this explains why the Calogero model in a harmonic background can be brought into the pure Calogero form [15]. Restoring the nonlinear term allows us to infer also that

$$i\partial_t\psi + \partial_x^2\psi + \left(-\frac{\omega^2 x^2}{4} + \frac{1}{\cos \omega t} |\psi|^2 \right) \psi = 0 \quad (21)$$

is integrable, and its solutions are obtained from those of the “free” NLS by the transformation (20). Let us mention that the covariance w. r. t. chronoprojective transformations was used before [16] for solving the NLS in oscillator and uniform-field backgrounds.

Now the constant-coefficient, damped, driven NLS,

$$i\partial_t\psi + \partial_x^2\psi + F|\psi|^2\psi = a(t, x)\psi + b(t, x), \quad (22)$$

passes the Painlevé test if

$$\begin{aligned} a(t, x) &= (\tfrac{1}{2}\partial_t\beta - \beta^2)x^2 + i\beta(t) + \alpha_1(t)x + \alpha_0(t), \\ b(t, x) &= 0 \end{aligned} \quad (23)$$

[17], i. e., precisely when the potential can be transformed away by our “non-relativistic conformal transformations”.

4 An integrable extension

Unfortunately, the time-dependent travelling soliton (10) is inconsistent with the original equation (5), since its phase is quadratic in x rather than linear, as required by consistency. The clue for finding integrable extensions is to observe that Eqn. (5) is in fact a Derivative Non-Linear Schrödinger equation (DNLS) [6]. Now the results of Clarkson and Cosgrove [18] say that the constant-coefficient equation

$$i\partial_t\psi + \frac{1}{2}\partial_x^2\psi + ia\psi\psi^*\partial_x\psi + ib\psi^2\partial_x\psi^* + c\psi^3\psi^* = 0 \quad (24)$$

is integrable iff

$$c = \frac{1}{2}b(2b - a). \quad (25)$$

In our case $a = -b = -\kappa^2$ and $c = 0$; Eq. (5) is therefore not integrable. However, adding a 6th-order potential to the Lagrange density i. e. considering rather

$$i\partial_t\psi + \frac{1}{2}\partial_x^2\psi + 2\kappa^2 j\psi + \frac{3}{2}\kappa^4|\psi|^4\psi = 0 \quad (26)$$

converts (5) into an integrable equation. Eq. (26) admits, e. g., the travelling wave solution $\psi = \sqrt{\rho}e^{i\theta}$, where

$$\rho = \frac{|v|}{2\kappa^2} \frac{1}{\sqrt{2} \cosh[v(x - \frac{v}{2}t)] + \text{sign } v}, \quad \theta = \frac{v}{2}x. \quad (27)$$

This can be checked by observing that for the Ansatz $\psi = f(x, t)e^{ivx/2}$ the modified NLS (26) again reduces to a constant-coefficient equation. Then the imaginary part of (26) requires that $f(x, t) = f(x - (v/2)t)$, while the real part can be integrated by the usual trick of multiplication by f' . The asymptotic conditions fix the integration constant to vanish, yielding a six-order non-linear equation, only containing even powers of f . Then, introducing $\rho = f^2$, we end up with the equation

$$(\rho')^2 - v^2\rho^2 + 4\kappa^2v\rho^3 + 4\kappa^4\rho^4 = 0, \quad (28)$$

whose integration provides us with (27).

Another way of understanding how the integrability comes is to apply the non-local transformation (4) backwards, which carries (26) into a Derivative Non-linear Schrödinger equation of type II (DNLSII),

$$i\partial_t\phi + \frac{1}{2}\partial_x^2\phi + 2i\kappa^2\rho\partial_x\phi = 0, \quad (29)$$

which, consistently with Eq. (25), is integrable [19].

References

- [1] S. C. Zhang, H. T. Hansson and S. Kivelson, *Phys. Rev. Lett.* **62**, 307 (1989).
- [2] D. Levi, L. Vinet and P. Winternitz, *Ann. Phys.* **230**, 101 (1994).
- [3] R. Jackiw and S.-Y. Pi, *Prog. Theor. Phys. Suppl.* **107**, 1 (1992); G. Dunne, *Self-dual Chern-Simons Theories*. Springer Lecture Notes in Physics (1995).
- [4] U. Aglietti, L. Griguolo, R. Jackiw, S.-Y. Pi and D. Seminara, *Phys. Rev. Lett.* **77**, 4406 (1996); R. Jackiw, *J. Non. Lin. Math. Phys.* **4**, 241 (1997).
- [5] S. J. Benetton Rabello, *Phys. Rev. Lett.* **76**, 4007 (1996).

- [6] H. Min and Q.-H. Park, *Phys. Lett.* **B243**, 621 (1996); L. Griguolo and D. Seminara, *Nucl. Phys.* **B516**, 467 (1998).
- [7] N. Joshi, *Phys. Lett.* **A 125**, 456-460 (1986).
- [8] J. Weiss, M. Tabor, and G. Carnevale, *J. Math. Phys.* **24**, 522 (1983).
- [9] P. A. M. Dirac, *Proc. R. Soc.* **A165**, 199 (1938);
- [10] J. P. Vinti, *Mon. Not. R. Astron. Soc.* **169**, 417 (1974).
- [11] C. Duval, G. Gibbons and P. A. Horváthy, *Phys. Rev.* **D43**, 3907 (1991).
- [12] R. Jackiw, *Physics Today*, **25**, 23 (1972); U. Niederer, *Helv. Phys. Acta* **45**, 802 (1972); C. R. Hagen, *Phys. Rev.* **D5**, 377 (1972). See also C. Duval, G. Burdet, H. P. Künzle and M. Perrin, *Phys. Rev.* **D31**, 1841 (1985).
- [13] H. H. Chen and C. S. Liu, *Phys. Rev. Lett.* **37**, 693 (1976); R. Balakrishnan, *Phys. Rev.* **A32**, 1144 (1985).
- [14] U. Niederer, *Helv. Phys. Acta* **46**, 192 (1973).
- [15] J. Avan and A. Jevicki, *Phys. Lett.* **B266**, 35 (1991).
- [16] G. Burdet and M. Perrin, *Lett. Math. Phys.* **11**, 147 (1986).
- [17] P. A. Clarkson, *Proc. Roy. Soc. Edin.* **A 109**, 109-126 (1988).
- [18] P. A. Clarkson and C. M. Cosgrove, *J. Phys. A: Math. Gen.* **20**, 2003 (1987).
- [19] D. J. Kaup and A. C. Newell, *Journ. Math. Phys.* **19**, 798 (1978); H. H. Chen, Y. C. Lee and C. S. Liu, *Physica Scripta* **20**, 490 (1979).